

Big impacts by small RNAs in plant development

George Chuck, Héctor Candela and Sarah Hake

The identification and study of small RNAs, including microRNAs and *trans*-acting small interfering RNAs, have added a layer of complexity to the many pathways that regulate plant development. These molecules, which function as negative regulators of gene expression, are now known to have greatly expanded roles in a variety of developmental processes affecting all major plant structures, including meristems, leaves, roots, and inflorescences. Mutants with specific developmental phenotypes have also advanced our knowledge of the biogenesis and mode of action of these diverse small RNAs. In addition, previous models on the cell autonomy of microRNAs may have to be revised as more data accumulate supporting their long distance transport. As many of these small RNAs appear to be conserved across different species, knowledge gained from one species is expected to have general application. However, a few surprising differences in small RNA function seem to exist between monocots and dicots regarding meristem initiation and sex determination. Integrating these unique functions into the overall scheme for plant growth will give a more complete picture of how they have evolved as unique developmental systems.

Addresses

Plant Gene Expression Center, United States Department of Agriculture-Agriculture Research Service and the University of California, Albany, CA 94710, United States

Corresponding author: Chuck, George (gchuck@nature.berkeley.edu)

Current Opinion in Plant Biology 2009, **12**:81–86

This review comes from a themed issue on
Growth and Development
Edited by Charles S. Gasser and Caroline Dean

Available online 6th November 2008

1369-5266/\$ – see front matter
Published by Elsevier Ltd.

DOI [10.1016/j.pbi.2008.09.008](https://doi.org/10.1016/j.pbi.2008.09.008)

Introduction

Since the last review in this journal of the role of microRNAs in development in plants [1], several advances in small RNA biology have been made, many novel and unexpected. This review focuses on how small RNAs have impacted our views of the development of several plant-specific structures in both monocots and dicots. While small RNAs have long been implicated in the determination of meristem boundaries in dicots, new data in monocots show that they are also essential for meristem initiation and maintenance. Small RNAs are known to

control the morphogenesis and abaxial/adaxial patterning of leaves, but also have been shown to affect the differentiation of specific cells types, such as the stomata, and the development of serrations. Two microRNAs known to control developmental timing have been shown to participate in a wider range of functions, including light response, control of plastochron length, and sex determination. The sequential expression patterns of this pair of plant-specific microRNAs are reminiscent of those of temporally regulated microRNAs in animals, indicating potential convergence of timing mechanisms between the two kingdoms. Finally, more evidence supporting the long distance transport and function of microRNAs has accumulated, a fact that will have to be incorporated into future models for microRNA action.

Meristems

In plants, organogenesis occurs from meristems, groups of self-organizing cells whose derivatives divide and elongate, making up the cells of the plant body. The root and shoot apical meristems form early in the embryo and produce opposite poles of growth and differentiation. Additional meristems such as lateral root meristems or axillary meristems form during the life of the plant, helping to create the overall architecture.

Shoot meristems have two major functions; they maintain a population of totipotent cells and produce lateral organs whose fates are determined by genetic and developmental factors. During early vegetative phases, the lateral organs are leaves that may have axillary meristems in the region between the leaf and the stem. In addition, leaves initiated during the juvenile vegetative phase have different morphology and cell identities compared to those made in the adult phase. Later, during reproductive development, the leaves are smaller or suppressed altogether, and the axillary meristems grow out into flowers or branches. Regardless of their size, leaves arise with an inherent polarity. The branches, on the other hand, are usually radially symmetric. Meristems are classified as determinate or indeterminate based on whether they are consumed in the production of primordia. Floral meristems are determinate since they terminate growth after the initiation of floral organs, while branch meristems are usually indeterminate.

Several interesting mutants have been described in rice that either lack or fail to properly maintain the shoot meristem. The *shootless* (*shl*) mutants initiate the coleoptile and scutellum, which together form the cotyledon in the grasses [2], but fail to make the shoot meristem [3]. Weak alleles show that *SHL* genes are also required for

maintenance of the meristem [4]. The *shoot meristem organization* (*sho*) mutants produce radialized leaves with irregular phyllotaxy [5], reminiscent of the maize mutant *leafbladeless1* (*lbl1*), which also produces radialized leaves with altered *KNOX* gene expression [6].

The *sho* and *shl* mutants of rice and the *lbl1* mutant of maize are defective in small RNA processing. *SHL2* encodes an RNA-dependent RNA polymerase 6, *SHO2* encodes an AGO7, *SHO1* encodes DCL4 [7^{••}], and *LBL1* encodes a SUPPRESSOR OF GENE SILENCING3 (SGS3) protein [8^{••}]. All of these proteins are components of the si-RNA pathway [9] and, interestingly, the mutant phenotypes are more severe in maize and rice than in *Arabidopsis*, where they mainly affect phase change. In the *sho*, *shl*, and *lbl1* mutants, the expression of class III *HD-ZIP* genes (*HD-ZIPIII*) is decreased, and the expression of the *MIR166* microRNA is increased [7^{••}, 8^{••}]. A critical role for these two components was shown by overexpressing a microRNA-resistant version of the *HD-ZIPIII* gene *OSHB1* in *sho1* mutants [7^{••}]. The phyllotaxy was restored to normal and the plants no longer produced radial leaves. Two other key components are likely to be the *AUXIN RESPONSE FACTOR* (*ARF*) genes and the *trans*-acting small interfering RNAs that target them, *tasiR-ARFs*. Levels of *tasiR-ARFs* are decreased in both *lbl1* mutants and *DCL4* knockdowns, with a corresponding increase in specific *ARF* target genes [8^{••}, 10[•]]. The unanswered question is how the regulation of *ARFs* by *tasi*-RNAs and *HD-ZIPIII* genes by *MIR166* are connected. Perhaps *MIR166* is positively regulated by *ARF* transcription factors, or, there exists an unknown *tasi*-RNA that negatively regulates the precursor of *MIR166*.

The *cup-shaped cotyledon* (*cuc*) mutant of *Arabidopsis* was identified by an embryo defect in which cotyledons are fused and leaves fail to initiate [11]. The redundant *CUC1* and *CUC2* genes are expressed in domains of the embryo that demarcate the position of the cotyledons, suggesting that they function to repress growth in that region [12, 13]. Because the *cuc1 cuc2* phenotype is seedling lethal and there is functional redundancy with *CUC3* [14, 15], the postembryonic role of *CUC* genes has not been well studied. A series of papers explores how the *MIR164* gene family modulates the expression of *CUC* genes and helps to tease out specific functions for them at different times.

Mutations in *MIR164c*, but not *MIR164b*, produce extra petals on the first formed flowers [16]. This mild phenotype is enhanced in the *MIR164a MIR164b MIR164c* triple mutant [17[•]]. In fact, most flowers of the triple mutant also have extra sepals, fewer stamens, and highly variable organ number and organ size. Floral organ number is also changed in plants that express a microRNA-resistant version of *CUC2* [18]. *CUC2* also participates in leaf de-

velopment, phyllotaxy, and axillary meristem development. Mutations in *MIR164a* as well as overexpression of microRNA-resistant versions of *CUC2* cause exaggerated leaf serrations. The serrations initiate as in wild-type but become enhanced due to growth repression by *CUC2* in the sinus region. Conversely, smooth margined leaves are found either with *MIR164* overexpression or in recessive *cuc2* mutants [18]. A role for *CUC* genes in the maintenance of phyllotaxy is seen in microRNA-resistant *CUC2* plants [19[•]]. Leaves are initiated correctly but, due to ectopic *CUC* expression, their growth is altered and thus the phyllotaxy is abnormal. This phenotype is also seen in the triple *MIR164a MIR164b MIR164c* mutant [17[•]]. Finally, reduction of *CUC2* and *CUC1* levels by overexpressing *MIR164* enhances a *cuc3* axillary meristem defect, revealing an additional role for *CUC1* and *CUC2* in this process [20].

Lateral organs — leaves

In *Arabidopsis*, *MIR165* and *MIR166* are two related microRNAs that differ at a single position but are thought to target the same set of *HD-ZIPIII* genes. In general, the overexpression of a microRNA is expected to recapitulate the loss-of-function phenotype of its targets. Now, Zhou *et al.* have found that overexpressing *MIR165* causes a phenotype different from that of *MIR166* overexpression. This phenotype matches the loss-of-function phenotype for *HD-ZIPIII* genes, an observation that may be explained if *MIR165* and *MIR166* target different genes with different efficiency [21]. If the many *HD-ZIPIII* proteins differ in their functions, understanding their expression patterns and the effects of their microRNA-resistant versions becomes particularly important. Itoh *et al.* [22] have carried out such a detailed study on the five members of the *HD-ZIPIII* family in rice, providing detailed expression data and setting the basis for additional functional studies.

Ori *et al.* have established the function of the *Lanceolate* (*LA*) gene of tomato in the morphogenesis of compound leaves [23^{••}]. Normal tomato leaves are compound and consist of several orders of leaflets, while semidominant *La* mutant leaves are simple. Cloning of the gene showed that *LA* encodes a TCP transcription factor, and that the semidominant alleles carry point mutations at a *MIR319*-binding site [23^{••}]. Interestingly, the expression domains of *LA* and *MIR319* partially overlap, suggesting that *MIR319* modulates, rather than fully downregulates, *LA* expression. Thus, the authors propose that *MIR319* and *LA* act together to regulate the competence to make leaflets, an idea supported by the simple leaves of *LA* overexpressors and the prolonged morphogenesis at the margins of *MIR319* overexpressors [23^{••}].

MicroRNAs not only affect leaf shape but also regulate cell differentiation within the leaf. A role for a microRNA, *MIR824*, in the patterning of stomatal complexes in

Arabidopsis was recently discovered [24]. *MIR824* targets the *MADS*-box gene *AGAMOUS-LIKE16* (*AGL16*), which is normally expressed in guard cells [25]. Plants expressing a *MIR824*-resistant version of *AGL16* develop higher order stomatal complexes as a result of the early formation of meristemoids, which takes place over a prolonged period. Consistent with this function, both *MIR824* and *AGL16* are expressed in the meristemoid lineage [24]. However, they do not occur simultaneously in the same cell types; while *MIR824* was expressed in the satellite meristemoids and guard mother cells, *AGL16* was expressed in differentiated guard cells.

Characterization of leaf mutants has continued to provide mechanistic insight into the biogenesis and mode of action of microRNAs. Analysis of a classic leaf mutant, *serrate* (*se*), has revealed an unexpected convergence between processing of microRNA precursors and protein-encoding transcripts. The serrated leaf phenotype of *se* mutants is similar to that of mutants of the cap-binding complex (CBC) involved in recruitment of splicing machinery. While *SE* was previously shown to function in primary microRNA processing [26,27], use of whole genome tiling arrays uncovered an additional function for *SE* in the splicing of normal transcripts [28]. This finding raises the possibility that *SE*, along with its interacting partners *DCL1* and *HYL1* may be part of a more extensive CBC complex, or may be a mediator between the microRNA-processing complex with the CBC [28]. Leaf serration similar to that of *se* mutants was also found in *abh1* single mutants, which reinforces the link with the miRNA pathway [29]. *ABH1* encodes a subunit of a nuclear CBC. Indeed, the primary transcripts of some microRNA genes as well as miRNA targets were represented at increased levels in the *abh1* single mutant. Gregory *et al.* hypothesize that ABH1 facilitates the correct processing of microRNA transcripts by binding to their 5'-ends and directly interacting with some protein of the miRNA biogenesis complex, perhaps *SE* [29].

Roots and plasmodesmata

While *MIR166* plays a major role in regulating *HD-ZIP* transcription factors controlling vascular development and polarity in the aerial portions of the plant, it also affects lateral root initiation. Work in *Medicago* showed that *MIR166* overexpression causes a decrease in lateral root number, and consequently a decrease in the number of nitrogen fixing root nodules [30]. Work on *MIR399*, which targets the *PHO2* gene that regulates phosphate homeostasis, revealed an intriguing case of long distance microRNA transport and function from shoots to roots. Grafts of shoots overexpressing *MIR399* onto wild-type roots showed accumulation of the mature microRNA and decreased *PHO2* transcript in root tissue [31,32]. These results suggest that microRNAs can function noncell autonomously, in contrast to previous studies showing that microRNAs function locally only where they are

expressed. Several microRNAs have been cloned from phloem sap in several species [33,34], but whether they are transported to the phloem through the cellular connections known as plasmodesmata (PD) is unknown. Recent work by Kobayashi *et al.* suggests a role for PD function in siRNA-mediated RNA silencing [35]. The *increased size exclusion limit2* (*ise2*) mutant alters PD morphology and continues to traffic large molecules at later stages of embryogenesis in contrast to normal embryos. Interestingly, *ise2* suppresses post-transcriptional gene silencing, although levels of microRNAs and siRNAs seem normal in the mutant. The *ISE2* gene encodes an RNA helicase, although it is unclear whether it regulates transcripts necessary for PD function or, like the pumpkin PSRP1 protein, it helps traffic small RNAs directly [34].

Flowering and sex determination

Work on the *MIR172* microRNA, which targets *APE-TALA2* (*AP2*) transcription factors, showed that it plays a major role in regulating floral meristem fate, organ identity, and flowering time [36,37]. A detailed study of *MIR172* function in flowers uncovered the mechanism behind its effects on floral meristem size and organ patterning. MicroRNA-resistant forms of *AP2* cause an enlarged indeterminate floral meristem, floral organ patterning defects, and extra whorls of stamens [36]. Zhao *et al.* showed that the floral meristem defect of these transformants is partially suppressed by *wuschel* (*wus*), but is synergistic with *agamous* (*ag*) [38], implicating both genes in the *AP2* pathway. In addition, the inner boundary of the *B*-class genes *APETALA3* and *PISTILLATA* was enlarged and shifted in the transformants, thus explaining the extra stamen defect. These data point to a more complex model for *AP2* function, where *AP2* does not simply repress the *C*-class gene *AG* as in the classic ABC model, but also acts in separate, parallel pathways to repress *WUS* and to define the boundary of *B*-class gene expression.

In *Petunia* and *Antirrhinum* a different microRNA, *MIR169*, seems to have adopted the role of *AP2*. The *blind* mutant of *Petunia* and the *fistulata* mutant of *Antirrhinum* show homeotic transformations in which second whorl organs are staminoid [39,40]. Both mutations are deletions of *MIR169* [41], which targets the *NF-YA* or CCAAT-box binding factor gene family [42]. These transcription factors are thought to regulate the *C*-class genes, whose expression changes in the *blind* and *fistulata* mutants [41].

MIR172 also plays a role in the photoperiodic flowering pathway in *Arabidopsis* and sex determination in maize. Earlier studies found that *MIR172* regulates flowering time through the repression of the *AP2* genes *TOE1* and *TOE2* [37], but it was not clear how *MIR172* responds to environmental signals to regulate these genes. Recent


studies have found that *MIR172* levels are decreased in several photoreceptor mutants, including *phyA*, *hy4*, *fha*, and *gigantea (gi)* [43]. *GI* encodes a nuclear protein involved in promoting flowering and regulating circadian rhythms. *MIR172* ultimately affects flowering time through regulation of the mobile floral inducer *FT*, which acts downstream of *TOE1* and *TOE2*, but in a pathway separate from *CONSTANS* [43]. Both *DCL1* and *SE* were also downregulated in the *gi* mutant, so it is possible that the late flowering phenotype is a result of an alteration in microRNA processing.

The involvement of *MIR172* in sex determination and meristem branching in maize inflorescences was discovered by the cloning of two classic maize mutants, the recessive *tasselseed4 (ts4)* mutant and the dominant *Tassel-seed6 (Ts6)* mutant. Both mutants belong to a class of *tasselseed* mutants that fail to abort pistils in the male inflorescence and increase indeterminate meristem branching in maize [44]. *TS4* encodes a unique microRNA of the *MIR172* family, *ZMA-MIR172E* [45^{*}]. A target of *TS4* is the *AP2* gene *INDETERMINATE SPIKELET1 (IDS1)*, whose mutations partially suppress the *ts4* phenotype. Cloning of *Ts6* revealed it to be the same gene as *IDS1*, except with mutations within the *MIR172*-binding site. While the meristem indeterminacy defects of *ts4* and *Ts6* are similar to those caused by microRNA-resistant forms of *AP2* in *Arabidopsis*, the sex determination defect is novel. *IDS1* and a related *ts4* target gene, *SISTER OF IDS1 (SID1)*, redundantly repress the maize *C*-class MADS-box genes *ZAG1* and *ZMM2* in bracts (in press). In *ts4* and *Ts6*, these MADS-box genes may be affected by the persistent expression of *AP2* genes, and somehow allow a bypass of the sex determination signal [45^{*}].

Timing and phase change

MicroRNAs have long been suspected of controlling the juvenile-to-adult vegetative phase transition in *Arabidopsis*, since many microRNA biogenesis mutants alter phase change [46]. In particular, the *MIR156* microRNA has been the subject of great scrutiny since the discovery that overexpression of microRNA-resistant versions of one of its target genes, *SPL3*, causes a truncation of the juvenile phase and early flowering [47]. Like *MIR172*, *MIR156* regulates these genes at the level of translation [48], which is proving to be a mode of microRNA regulation in plants more common than previously thought [49]. In addition, overexpression of *MIR156* causes late flowering and extension of the juvenile phase [47,50]. In maize, a similar phenotype is seen in the dominant mutant *Corn-grass1 (Cg1)* [51^{*}]. Cloning of *Cg1* revealed that it encodes a tandem *MIR156* gene that is overexpressed in the mutant. Interestingly, another *MIR172* target, *GLOSSY15*, which maintains juvenile leaf cell identities, is ectopically expressed in the *Cg1* mutant [52]. This observation may be explained by the presence of reduced levels of *MIR172* in *Cg1* leaves that overexpress *MIR156*

Figure 1



Reciprocal expression patterns of *MIR156* and *MIR172* in the juvenile and adult phase of development.

[51^{*}]. Thus, there appears to be a converse regulatory relationship between *MIR156* and *MIR172*, where the relative balance of each may be important for the timing of the phase transition (Figure 1).

Other targets of *MIR156* play different roles in shoot development in *Arabidopsis*. For example, *MIR156* overexpression causes the overproduction of leaves and a shortened plastochron [47,50]. These phenotypes are primarily due to downregulation of *SPL9* and *SPL15* [53], which act to limit leaf initiation rates. By using tissue-specific promoters to drive *MIR156* expression, it was shown that the shortened plastochron effect could not be induced with meristem-specific expression, but only with lateral organ expression [54], indicating a noncell autonomous effect of either *MIR156* or one of its *SPL* targets on the meristem. The *SPL9* gene represents a likely noncell autonomous target since it is not expressed in the meristem, and yet can affect plastochron when *MIR156* resistant forms are expressed in either leaf primordia or the meristem [54].

There are several interesting parallels between plant and animals in the control of stage-specific developmental events by microRNAs. In *C. elegans* the transition from early larval stages to the mature adult is controlled by the sequential expression of the *lin-4* and *let-7* microRNAs [55], the functions of which appear to be conserved in other animal species [56]. Along the same lines, the transition from juvenile to adult shoot development in plants appears to be regulated by the *MIR156* and *MIR172* microRNAs, both of which appear to be sequentially expressed and conversely regulated. As in animals, both microRNAs control genes that specify cell identities specific to each stage of development. In maize, *MIR156* appears to be

present during the juvenile phase [51[•]] when *MIR172* is absent, and the converse is true in the adult [45[•],57] (Figure 1). These reciprocal expression patterns of *MIR156* and *MIR172* appears to be conserved in *Arabidopsis* as well [37,47]. It will be critical to examine this converse relationship in other plants, and, more importantly, to associate different juvenile-to-adult transition times with the gain or loss of each microRNA. Also intriguing is the fact that both these microRNAs were found in phloem sap [33], opening the possibility that they also act noncell autonomously to influence developmental timing.

Conclusion

This review of small RNA function in plant development provides just a snapshot of the many pathways affected by these small molecules. Future work on the transcriptional regulation of these microRNAs will give a clearer picture of how they are able to regulate their targets in such a precise manner. For example, what factors are responsible for turning *MIR156* off and *MIR172* on during the transition to the adult phase? Also, the evidence that microRNAs can move through graft junctions and affect gene expression at a distance provides the implication, but not the proof, that they may act noncell autonomously. Is this a general theme for all small RNAs? As more small RNAs become sequenced from different organisms there is little doubt that many other novel processes will be found to be under the control of these multifaceted molecules.

Acknowledgements

This work was supported by DOE grant DE-A102-08ER15962. We thank China Lunde for helpful comments on the manuscript.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Willmann M, Poethig RS: **Conservation and evolution of miRNA regulatory programs in plant development.** *Curr Opin Plant Biol* 2007, **10**:503-511.
2. Kaplan DR: **The monocotyledons: their evolution and comparative biology. VII. The problem of leaf morphology and evolution in the monocotyledons.** *Q Rev Biol* 1973, **48**:437-457.
3. Satoh N, Hong SK, Nishimura A, Matsuoka M, Kitano H, Nagato Y: **Initiation of shoot apical meristem in rice: characterization of four SHOOTLESS genes.** *Development* 1999, **126**:3629-3636.
4. Satoh N, Itoh J, Nagato Y: **The SHOOTLESS2 and SHOOTLESS1 genes are involved in both initiation and maintenance of the shoot apical meristem through regulating the number of indeterminate cells.** *Genetics* 2003, **164**:335-346.
5. Itoh JI, Kitano H, Matsuoka M, Nagato Y: **Shoot organization genes regulate shoot apical meristem organization and the pattern of leaf primordium initiation in rice.** *Plant Cell* 2000, **12**:2161-2174.
6. Timmermans MC, Schultes NP, Jankovsky JP, Nelson T: **leafbladeless1 is required for dorsoventrality of lateral organs in maize.** *Development* 1998, **125**:2813-2823.
7. Nagasaki H, Itoh J, Hayashi K, Hibara K, Satoh-Nagasawa N, •• Nosaka M, Mukouhata M, Ashikari M, Kitano H, Matsuoka M et al.: **The small interfering RNA production pathway is required for shoot meristem initiation in rice.** *Proc Natl Acad Sci U S A* 2007, **104**:14867-14871.
- Three different shoot meristem mutants were cloned by position and shown to function in the siRNA pathway. Both tasiRNA and *mir166* are implicated in meristem initiation and maintenance.
8. Nogueira F, Madi S, Chitwood D, Juarez MT, Timmermans MC: •• **Two small regulatory RNAs establish opposing fates of a developmental axis.** *Genes Dev* 2007, **21**:750-755.
- The *leafbladeless1* mutation was cloned and shown to encode SGS3. An interaction between the small interfering RNA pathway and *mir166* is demonstrated.
9. Mallory ACBN: **MicroRNA-directed regulation: to cleave or not to cleave.** *Trends Plant Sci* 2008, **13**:359-367.
10. Liu B, Chen Z, Song X, Liu C, Cui X, Zhao X, Fang J, Xu W, Zhang H, • Wang X et al.: **Oryza sativa dicer-like4 reveals a key role for small interfering RNA silencing in plant development.** *Plant Cell* 2007, **9**:2705-2718.
- The authors analyze recessive *dcl4* mutants in rice and observe radialized floral organs. Levels of siRNAs were affected in the *dcl4* mutant.
11. Aida M, Ishida T, Fukaki H, Fujisawa H, Tasaka M: **Genes involved in organ separation in Arabidopsis: an analysis of the cup-shaped cotyledon mutant.** *Plant Cell* 1997, **9**:841-857.
12. Aida M, Ishida T, Tasaka M: **Shoot apical meristem and cotyledon formation during Arabidopsis embryogenesis: interaction among the CUP-SHAPED COTYLEDON and SHOOT MERISTEMLESS genes.** *Development* 1999, **126**:1563-1570.
13. Takada S, Hibara K, Ishida T, Tasaka M: **The CUP-SHAPED COTYLEDON1 gene of Arabidopsis regulates shoot apical meristem formation.** *Development* 2001, **128**:1127-1135.
14. Vroemen CW, Mordhorst AP, Albrecht C, Kwaaitaal MA, de Vries SC: **The CUP-SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in Arabidopsis.** *Plant Cell* 2003, **15**:1563-1577.
15. Hibara K, Takada S, Tasaka M: **CUC1 gene activates the expression of SAM-related genes to induce adventitious shoot formation.** *Plant J* 2003, **36**:687-696.
16. Baker CC, Sieber P, Wellmer F, Meyerowitz EM: **The early extra petals1 mutant uncovers a role for microRNA mir164c in regulating petal number in Arabidopsis.** *Curr Biol* 2005, **15**:303-315.
17. Sieber P, Wellmer F, Gheyselsinck J, Riechmann JL, • Meyerowitz EM: **Redundancy and specialization among plant microRNAs: role of the MIR164 family in developmental robustness.** *Development* 2007, **134**:1051-1060.
- The phenotype of *MIR164* triple mutants is analyzed as well as the specific expression patterns. *CUC2* RNA levels are increased in the triple mutant, but the domain of expression does not change.
18. Nikovics K, Blein T, Peaucelle A, Ishida T, Morin H, Aida M, Laufs P: **The balance between the MIR164A and CUC2 genes controls leaf margin serration in Arabidopsis.** *Plant Cell* 2006, **18**:2929-2945.
19. Peaucelle A, Morin H, Traas J, Laufs P: **Plants expressing a MIR164-resistant CUC2 gene reveal the importance of post-meristematic maintenance of phyllotaxy in Arabidopsis.** *Development* 2007, **134**:1045-1050.
- The authors carefully analyze the phyllotaxy defect of microRNA-resistant *CUC2* plants.
20. Raman S, Greb T, Peaucelle A, Blein T, Laufs P, Theres K: **Interplay of MIR164, CUP-SHAPED COTYLEDON genes and LATERAL SUPPRESSOR controls axillary meristem formation in Arabidopsis thaliana.** *Plant J* 2008, **55**:65-76.
21. Zhou GK, Kubo M, Zhong R, Demura T, Ye ZH: **Overexpression of MIR165 affects apical meristem formation, organ polarity establishment and vascular development in Arabidopsis.** *Plant Cell Physiol* 2007, **48**:391-404.
22. Itoh JI, Hibara KI, Sato Y, Nagato Y: **Developmental role and auxin responsiveness of Class III HD-Zip gene family members in rice.** *Plant Physiol* 2008, **147**:1960-1975.

23. Ori N, Cohen AR, Etzioni A, Brand A, Yanai O, Shleizer S, Menda N, Amsellem Z, Efroni I, Pekker I *et al.*: **Regulation of LANCEOLATE by MIR319 is required for compound-leaf development in tomato.** *Nat Genet* 2007, **39**:941-951.
- The authors identify the gene responsible for the *Lanceolate* phenotype, a classical mutant of tomato, and show that a microRNA regulates the competence to make compound leaves.
24. Kutter C, Schöb H, Stadler M, Meins F, Si-Ammour A: **MicroRNA-mediated regulation of stomatal development in Arabidopsis.** *Plant Cell* 2007, **19**:2417-2429.
25. Alvarez-Buylla ER, Liljegren SJ, Pelaz S, Gold SE, Burgeff C, Ditta GS, Vergara-Silva F, Yanofsky MF: **MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes.** *Plant J* 2000, **24**:457-466.
26. Lobbes D, Rallapalli G, Schmidt DD, Martin C, Clarke J: **SERRATE: a new player on the plant microRNA scene.** *EMBO J* 2006, **25**:1052-1058.
27. Yang L, Liu Z, Dong A, Huang H: **SERRATE is a novel nuclear regulator in primary microRNA processing in Arabidopsis.** *Plant J* 2006, **47**:841-850.
28. Laubinger S, Sachsberg T, Zeller G, Busch W, Lohmann JU, Ratsch G, Weigel D: **Dual roles of the nuclear cap-binding complex and SERRATE in pre-mRNA splicing and microRNA processing in Arabidopsis thaliana.** *Proc Natl Acad Sci U S A* 2008, **105**:8795-8800.
29. Gregory BD, O'Malley RC, Lister R, Ulrich MA, Tonti-Filippini J, Chen H, Millar AH, Ecker JR: **A link between RNA metabolism and silencing affecting Arabidopsis development.** *Dev Cell* 2008, **14**:854-866.
30. Boualem A, Laporte P, Jovanovic M, Laffont C, Plet J, Combier JP, Niebel A, Crespi M, Frugier F: **MicroRNA166 controls root and nodule development in Medicago truncatula.** *Plant J* 2008, **54**:876-887.
31. Pant BD, Buhtz A, Kehr J, Scheible WR: **MicroRNA399 is a long-distance signal for the regulation of plant phosphate homeostasis.** *Plant J* 2008, **53**:731-738.
32. Lin SI, Chiang SF, Lin WY, Chen JW, Tseng CY, Wu PC, Chiou TJ: **Regulatory network of microRNA399 and PHO2 by systemic signaling.** *Plant Physiol* 2008, **147**:732-746.
- Both Pant *et al.* and Lin *et al.* performed grafts of *MIR399* overexpressing shoots to wild-type roots and found an increase of mature *MIR399* and a reduction of *PHO2* transcript in roots. This is a unique example of microRNAs affecting gene expression at long distances.
33. Buhtz A, Springer F, Chappell L, Baulcombe DC, Kehr J: **Identification and characterization of small RNAs from the phloem of Brassica napus.** *Plant J* 2008, **53**:739-749.
34. Yoo BC, Kragler F, Varkonyi-Gasic E, Haywood V, Archer-Evans S, Lee YM, Lough TJ, Lucas WJ: **A systemic small RNA signaling system in plants.** *Plant Cell* 2004, **16**:1979-2000.
35. Kobayashi K, Otegui MS, Krishnakumar S, Mindrinos M, Zambryski P: **INCREASED SIZE EXCLUSION LIMIT 2 encodes a putative DEVH box RNA helicase involved in plasmodesmata function during Arabidopsis embryogenesis.** *Plant Cell* 2007, **19**:1885-1897.
36. Chen X: **A microRNA as a translational repressor of APETALA2 in Arabidopsis flower development.** *Science* 2004, **303**:2022-2025.
37. Aukerman MJ, Sakai H: **Regulation of flowering time and floral organ identity by a microRNA and its APETALA2-like target genes.** *Plant Cell* 2003, **15**:2730-2741.
38. Zhao L, Kim Y, Dinh TT, Chen X: **MIR172 regulates stem cell fate and defines the inner boundary of APETALA3 and PISTILLATA expression domain in Arabidopsis floral meristems.** *Plant J* 2007, **51**:840-849.
- A complex pathway for *AP2* gene function is proposed that acts through the *WUSCHEL* and *AGAMOUS* pathways to affect floral meristem size, determinacy, and organ identity.
39. McSteen PC, Vincent CA, Doyle S, Carpenter R, Coen ES: **Control of floral homeotic gene expression and organ morphogenesis in Antirrhinum.** *Development* 1998, **125**:2359-2369.
40. Motte P, Saedler H, Schwarz-Sommer Z: **STYLOSA and FISTULATA: regulatory components of the homeotic control of Antirrhinum floral organogenesis.** *Development* 1998, **125**:71-84.
41. Cartolano MCR, Efremova N, Kuckenberg M, Zethof J, Gerats T, Schwarz-Sommer Z, Vandenbussche M: **A conserved microRNA module exerts homeotic control over Petunia hybrida and Antirrhinum majus floral organ identity.** *Nat Genet* 2007, **39**:901-905.
- Two naturally occurring mutants in *Petunia* and *Antirrhinum* are shown to be due to mutations in *MIR169* which appears to prevent ectopic expression of class C genes in these species.
42. Jones-Rhoades MW, Bartel DP, Bartel B: **MicroRNAs and their regulatory roles in plants.** *Annu Rev Plant Biol* 2006, **57**:19-53.
43. Jung JH, Seo YH, Seo PJ, Reyes JL, Yun J, Chua NH, Park CM: **The GIGANTEA-regulated microRNA172 mediates photoperiodic flowering independent of CONSTANS in Arabidopsis.** *Plant Cell* 2007, **19**:2736-2748.
44. Irish EE: **Class II tassel seed mutations provide evidence for multiple types of inflorescence meristems in maize (Poaceae).** *Am J Bot* 1997, **84**:1502-1515.
45. Chuck G, Meeley RB, Irish E, Sakai H, Hake S: **The maize tasselseed4 microRNA controls sex determination and meristem cell fate by targeting Tasselseed6/indeterminate spikelet1.** *Nat Genet* 2007, **39**:1517-1521.
- MIR172* and target genes in maize are shown to control sex determination and branching in the inflorescence.
46. Willmann MR, Poethig RS: **Time to grow up: the temporal role of small RNAs in plants.** *Curr Opin Plant Biol* 2005, **8**:548-552.
47. Wu G, Poethig RS: **Temporal regulation of shoot development in Arabidopsis thaliana by MIR156 and its target SPL3.** *Development* 2006, **133**:3539-3547.
48. Gandikota M, Birkenbihl RP, Hohmann S, Cardon GH, Saedler H, Huijser P: **The miRNA156/157 recognition element in the 3'UTR of the Arabidopsis SBP box gene SPL3 prevents early flowering by translational inhibition in seedlings.** *Plant J* 2007, **49**:683-693.
49. Brodersen P, Sakvarelidze-Achard L, Bruun-Rasmussen M, Dunoyer P, Yamamoto YY, Sieburth L, Voinnet O: **Widespread translational inhibition by plant miRNAs and siRNAs.** *Science* 2008, **320**:1185-1190.
50. Schwab R, Palatnik JF, Rießer M, Schommer C, Schmid M, Weigel D: **Specific effects of microRNAs on the plant transcriptome.** *Dev Cell* 2005, **8**:517-527.
51. Chuck G, Cigan M, Saetern K, Hake S: **The heterochronic maize mutant Corngrass1 results from overexpression of a tandem microRNA.** *Nat Genet* 2007, **39**:544-549.
- The pleiotropic *Cg1* mutant is shown to be due to overexpression of *MIR156* with a concomitant decrease in *MIR172*.
52. Moose SP, Sisco PH: **glossy15, an APETALA2-like gene from maize that regulates leaf epidermal cell identity.** *Genes Dev* 1996, **10**:3018-3027.
53. Schwarz S, Grande AV, Bujdosó N, Saedler H, Huijser P: **The microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in Arabidopsis.** *Plant Mol Biol* 2008, **67**:183-195.
54. Wang JW, Schwab R, Czech B, Mica E, Weigel D: **Dual effects of MIR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size in Arabidopsis thaliana.** *Plant Cell* 2008, **20**:1231-1243.
55. Moss EG: **Heterochronic genes and the nature of developmental time.** *Curr Biol* 2007, **17**:425-434.
56. Pasquinelli AE, Ruvkun G: **Control of developmental timing by microRNAs and their targets.** *Annu Rev Cell Dev Biol* 2002, **18**:495-513.
57. Lauter N, Kampani A, Carlson S, Goebel M, Moose SP: **microRNA172 down-regulates glossy15 to promote vegetative phase change in maize.** *Proc Natl Acad Sci U S A* 2005, **102**:9412-9417.